

Target Corporation

-PE Product Labels –

**SAFETY SEALED: DO NOT USE IF THE INDIVIDUAL
BLISTER UNIT IS OPEN OR TORN**

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms: nasal congestion, headache, cough, minor aches & pains, sore throat
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 softgels in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blistering
- Rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have liver disease, heart disease, diabetes, high blood pressure, thyroid disease, trouble urinating due to enlarged prostate gland, persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, or cough that occurs with too much phlegm (mucus).

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if nervousness, dizziness, or sleeplessness occur; pain, nasal congestion, or cough gets worse or lasts more than 7 days; fever gets worse or lasts more than 3 days; redness or swelling is present; new symptoms occur; or cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Other information

- Store at 20-25°C (68-77°F)
- Avoid excessive heat.

Inactive ingredients: FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide.

Compare to active ingredients in
Mucinex® Fast-Max® Cold & Flu*

maximum strength
cold and flu

acetaminophen
dextromethorphan HBr
guaifenesin
phenylephrine HCl

fast dissolving softgels

relieves aches, fever and
sore throat controls cough
relieves nasal and chest
congestion thins and
loosens mucus

up&up

16 SOFTGELS

ACTUAL SIZE

AGES
12+
YEARS

Varnish Omit Area

Varnish Omit Area

Print/Varnish Omit Area

Print/Varnish Omit Area

Lot #:

Exp. Date:

Questions? Call 1-877-290-4008

Distributed by Target Corporation
Minneapolis, MN 55403

Made in India TM & ©2020 Target Brands, Inc.

*This product is not manufactured or distributed by Reckitt Benckiser LLC, owner of the registered trademark Mucinex® Fast-Max® Cold & Flu. ©2017 0520

PAPER BOX

RECYCLE

TRAY

094 14 0077 R01 C-001227-01-055

3 49483 628161 6





Children's Day Time Cold + Cough Drug Facts

Drug Facts (continued)

Directions

● do not take more than 6 doses in any 24-hour period

● measure only with dosing cup provided. Do not use any other dosing device

● keep dosing cup with product

● mL = milliliter

Age	Dose
adults and children 12 years and over	20 mL, every 4 hours
children 6 to 11 years	10 mL, every 4 hours
children under 6 years	do not use

Other information

● each 10 mL contains sodium 1 mg

● store between 20°-25° (68°-77°). Do not refrigerate.

Inactive ingredients

citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucrose

Questions or comments?

Call 1-800-845-6024

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetapp® Cold & Cough.

100% satisfaction guaranteed or your money back.

Distributed by Target Corporation
Minneapolis, MN 55403
©2017 Target Brands, Inc.

Children's Night Time Cold + Congestion Drug Facts

Drug Facts

Active ingredients (in each 10 mL)

Diphenhydramine HCl 12.5 mg Antihistamine
Dextromethorphan HBr 10 mg Cough suppressant
Phenylephrine HCl 5 mg Nasal decongestant

Purposes

● temporarily relieves these symptoms occurring with a cold, hay fever, or other upper respiratory allergies

● nasal congestion ● runny nose ● sneezing

● itchy, watery eyes ● itching of the nose or throat

Warnings

Do not use

● to sedate a child or to make a child sleep

● if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product, or with any other product containing diphenhydramine, even one used on days

Ask a doctor before use if you have

● heart disease ● high blood pressure ● thyroid disease

● diabetes ● glaucoma

● trouble urinating due to an enlarged prostate gland

● cough that occurs with too much phlegm (mucus)

● a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

● taking any other oral nasal decongestant or stimulant

● taking sedatives or tranquilizers

When using this product

● do not use more than directed

● marked drowsiness may occur

● avoid alcoholic drinks

● alcohol, sedatives, and tranquilizers may increase drowsiness

● be careful when driving a motor vehicle or operating machinery or machinery may occur, especially in children

Stop use and ask a doctor if

● you get nervous, dizzy, or sleepless

● symptoms do not get better within 7 days or occur with a fever

● cough lasts more than 7 days, comes back, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Children's Night Time Cold + Congestion Drug Facts

Drug Facts (continued)

Directions

● do not take more than 6 doses in any 24-hour period

● do not exceed recommended dosage

● measure only with dosing cup provided. Do not use any other dosing device

● keep dosing cup with product

● mL = milliliter

Age	Dose
adults and children 12 years and over	20 mL, every 4 hours
children 6 to 11 years	10 mL, every 4 hours
children under 6 years	do not use

Other information

● each 10 mL contains sodium 0 mg

● store between 20°-25° (68°-77°)

● do not refrigerate

● protect from light

Inactive ingredients

sucrose, potassium, erythrasin, citric acid, FD&C blue #1, FD&C red #40, flavor, methyl, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol

Questions or comments?

Call 1-800-845-6024

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetapp® Nighttime Cold & Congestion.

100% satisfaction guaranteed or your money back.

Distributed by Target Corporation
Minneapolis, MN 55403
©2017 Target Brands, Inc.

Compare to active ingredients in Children's Dimetapp® Cold & Cough*

children's day time cold + cough

brompheniramine maleate 2 mg (antihistamine)

dextromethorphan HBr 10 mg (cough suppressant)

phenylephrine HCl 5 mg (nasal decongestant)

cough

itchy, watery eyes

runny nose

sneezing

stuffy nose

itching of the nose or throat

alcohol free

up & up

4 FL OZ (118 mL)

8 FL OZ (236 mL) TOTAL

GRAPE FLAVOR

DOSING CUP INCLUDED

AGES 6+ YEARS

NDC 11673-059-06

Compare to active ingredients in Children's Dimetapp® Nighttime Cold & Congestion**

children's night time cold + congestion

diphenhydramine HCl 12.5 mg (antihistamine/cough suppressant)

phenylephrine HCl 5 mg (nasal decongestant)

stuffy nose

runny nose

sneezing

itchy, watery eyes

cough

GRAPE FLAVOR

DOSING CUP INCLUDED

AGES 6+ YEARS

084 14 0026 R01

100-000121-01-153



7 15255 33608 6

PLD-82810 PZD-01178

PLD-8275A

Lot No.:

3084

nighttime cold and flu (continued)	daytime severe cold	daytime severe cold (continued)	NDC 11673-518-24	nighttime cold and flu	
Drug Facts (continued) occurs with smoking, asthma, chronic bronchitis, or emphysema cough that occurs with too much phlegm (mucus) Ask a doctor or pharmacist before use if you are: taking the blood thinning drug warfarin taking antacids or tranquilizers When using this product: do not use more than directed excitability may occur, especially in children marked drowsiness may occur alcohol, antacids, and tranquilizers may increase drowsiness be careful when driving a motor vehicle or operating machinery Stop use and ask a doctor if: numbness, dizziness, or sleeplessness occur pain, nasal congestion, or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-232-1223) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions do not take more than directed (see Overdose warning) do not take more than 12 softgels (Daytime and Nighttime) in any 24-hour period adults and children 12 years of age and older: take 2 softgels every 4 hours children under 12 years of age: do not use when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing. Other information swallow whole; do not crush, chew, or dissolve store between 15-30°C (59-86°F) avoid excessive heat Inactive ingredients HCl, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, stearic acid, sodium hydroxide, sorbitol, sorbitol, titanium dioxide	Drug Facts Active ingredients (in each softgel) Acetaminophen 325 mg Pain reliever/fever reducer Dextromethorphan HBr 10 mg Cough suppressant Guaifenesin 200 mg Expectorant Phenylephrine HCl 5 mg Nasal decongestant Uses temporarily relieves these common cold and flu symptoms headache nasal congestion sore throat cough minor aches and pains helps loosen phlegm (mucus) and the bronchial secretions to let the bronchial passageways of both nose and mouth and make coughs more productive temporarily reduces fever Warnings Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness blisters rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, hoarse, or vomiting, consult a doctor promptly. Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription medication for seizure (anticonvulsant) or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have: liver disease diabetes high blood pressure heart disease thyroid disease trouble urinating due to an enlarged prostate gland a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema cough that occurs with too much phlegm (mucus) Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.	Drug Facts (continued) When using this product, do not use more than directed. Stop use and ask a doctor if: numbness, dizziness, or sleeplessness occur pain, nasal congestion, or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-232-1223) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions do not take more than directed (see Overdose warning) do not take more than 12 softgels (Daytime and Nighttime) in any 24-hour period adults and children 12 years of age and older: take 2 softgels every 4 hours children under 12 years of age: do not use when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing. Other information swallow whole; do not crush, chew, or dissolve store between 15-30°C (59-86°F) avoid excessive heat Inactive ingredients FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, titanium dioxide Questions? Call 1-800-980-8874 *This product is not manufactured or distributed by Reichle Bionics, distributor of Maximum Strength Mucinex® Fast-Max® Day Severe Cold and Maximum Strength Mucinex® Fast-Max® Night Cold & Flu.	Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Day Severe Cold* maximum strength daytime severe cold acetaminophen (pain reliever/fever reducer) dextromethorphan HBr (cough suppressant) guaifenesin (expectorant) phenylephrine HCl (nasal decongestant) relieves aches, fever and sore throat controls cough relieves nasal and chest congestion thins and loosens mucus  ACTUAL SIZE AGES 12+ YEARS	Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Night Cold & Flu* maximum strength nighttime cold and flu acetaminophen (pain reliever/fever reducer) dextromethorphan HBr (cough suppressant) dextromethorphan HBr (cough suppressant) doxylamine succinate (antihistamine) phenylephrine HCl (nasal decongestant) relieves aches, fever and sore throat controls cough relieves nasal congestion relieves runny nose and sneezing  ACTUAL SIZE AGES 12+ YEARS	Drug Facts Active ingredients (in each softgel) Acetaminophen 325 mg Pain reliever/fever reducer Dextromethorphan HBr 10 mg Cough suppressant Doxylamine succinate 6.25 mg Antihistamine Phenylephrine HCl 5 mg Nasal decongestant Uses temporarily relieves these common cold and flu symptoms headache nasal congestion sore throat minor aches and pains runny nose and sneezing controls cough to help you get to sleep temporarily reduces fever Warnings Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness blisters rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, hoarse, or vomiting, consult a doctor promptly. Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription medication for seizure (anticonvulsant) or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have: liver disease high blood pressure diabetes heart disease thyroid disease glucose trouble urinating due to an enlarged prostate gland a breathing problem or chronic cough that lasts or as

094 14 0126 R00 C-000703-01-089



3 59726 51824 6

 Distributed by Target Corporation
 Minneapolis, MN 55403
 TM & ©2018 Target Brands, Inc.

PLU-A4218 F0204221

PARENTS:

www.Target.com/parents

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BULSTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

 Lot / Exp:
 Product of:

3085

nighttime cold and flu (continued)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.**Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- swallow whole; do not crush, chew, or dissolve
- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients: D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol*, polyethylene glycol, povidone, propylene glycol, purified water, shellac*, sodium hydroxide*, sorbitan, sorbitol, titanium dioxide

*contains one or more of these ingredients

Questions or comments?

Call 1-800-910-0874

daytime severe cold

Drug Facts

Active ingredients
(in each softgel)

Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant

Purposes

- temporarily relieves these common cold and flu symptoms
- headache
- nasal congestion
- sore throat
- cough
- minor aches and pains
- helps loosen phlegm (mucus) and the bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- high blood pressure
- heart disease
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor before use if you have

- liver disease
- diabetes
- high blood pressure
- heart disease
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

daytime severe cold (continued)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

- Stop use and ask a doctor if
- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels (Daytime and Nighttime) in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- swallow whole; do not crush, chew, or dissolve
- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients: FD&C red #40, FD&C yellow #6, gelatin, glycerin, mannitol*, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide

*may contain this ingredient

Questions or comments?

Call 1-800-910-0874

***This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Day Severe Cold and Maximum Strength Mucinex® Fast-Max® Night Cold & Flu.

Compare to active ingredients in
**Maximum Strength Mucinex®
Fast-Max® Day Severe Cold*****

maximum strength daytime severe cold

acetaminophen

(pain reliever/fever reducer)

dextromethorphan HBr

(cough suppressant)

guaifenesin (expectorant)

phenylephrine HCl

(nasal decongestant)

relieves aches, fever and sore throat
controls cough
relieves nasal and chest congestion
thins and loosens mucus



ACTUAL SIZE

AGES

12+

YEARS

16 SOFTGELS** (**LIQUID-FILLED CAPSULES)
24 TOTAL SOFTGELS

NDC 11673-920-24

nighttime cold and flu

Drug Facts

Active ingredients
(in each softgel)

Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Doxylamine succinate 6.25 mg.....Antihistamine
Phenylephrine HCl 5 mg.....Nasal decongestant

Purposes

- temporarily relieves these common cold and flu symptoms
- cough
- headache
- minor aches and pains
- sore throat
- nasal congestion
- runny nose and sneezing
- controls cough to help you get to sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- high blood pressure
- heart disease
- glaucoma
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor before use if you have

- liver disease
- diabetes
- high blood pressure
- heart disease
- glaucoma
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

094 14 0126 R01 C-000703-01-089



3 59726 51824 6

PARENTS:

Learn about how medicine works

www.StopMedicineAbuse.org

Distributed by Target Corporation
Minneapolis, MN 55403
TM & ©2019 Target Brands, Inc.

PLD-9421C; F0305488

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot / Exp.:
Product of:



No print/No varnish area
Adhesive Flap

No print/No varnish area
Adhesive

Drug Facts
KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT INFORMATION

Active ingredients (in each caplet)

Acetaminophen 325 mg.....Pain reliever

Guaifenesin 200 mg.....Expectorant

Phenylephrine HCl 5 mg.....Nasal decongestant

Active ingredients (in each caplet)

Acetaminophen 325 mg.....Pain reliever

Diphenhydramine HCl 12.5 mg.....Antihistamine/cough suppressant

Phenylephrine HCl 5 mg.....Nasal decongestant

Uses ■ temporarily relieve:
■ nasal congestion ■ headache ■ minor aches and pains
■ sinus congestion and pressure ■ cough (nighttime only)
■ runny nose and sneezing (nighttime only)
■ temporarily promotes nasal and/or sinus drainage
■ helps loosen phlegm (mucus) and thin bronchial secretions to aid the
bronchial passages of bothersome mucus and make coughs
more productive (daytime only)

Warnings
Liver warning: This product contains acetaminophen. Severe liver
damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions.
Symptoms may include: ■ skin redness ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Do not use
■ with any other drug containing acetaminophen (prescription or
nonprescription). If you are not sure whether a drug contains
acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor
(MAOI) (certain drugs for depression, psychiatric or emotional
disorders, or Parkinson's disease) or for 2 weeks after stopping
such a drug (if you or your doctor or pharmacist believe taking this
product, an MAOI, ask a doctor or pharmacist before taking this
product.
■ if you have ever had an allergic reaction to this product or any of its
ingredients

PEEL HERE FOR MORE DRUG FACTS

No print/No varnish area
Adhesive

No print/No varnish area
Lot no/Exp date

No print/No varnish area
Adhesive

No print/No varnish area
Lot no/Exp date

ADHESIVE AREA

Drug Facts (continued)

■ with any other product containing diphenhydramine, even one
used on skin (Nighttime only)

Ask a doctor before use if you have

■ heart disease ■ liver disease ■ diabetes
■ thyroid disease ■ high blood pressure
■ difficulty in urination due to enlargement of the prostate gland
■ persistent or chronic cough such as occurs with smoking,
asthma, chronic bronchitis, or emphysema
■ cough that occurs with too much phlegm (mucus)
■ a breathing problem such as emphysema or chronic bronchitis
(Nighttime only) ■ glaucoma (Nighttime only)

Ask a doctor or pharmacist before use if you are

■ taking the blood thinning drug warfarin
■ taking sedatives or tranquilizers (Nighttime only)

When using this product

■ do not exceed recommended dosage
■ excitability may occur, especially in children (Nighttime only)
■ marked drowsiness may occur (Nighttime only)
■ alcohol, sedatives, and tranquilizers may increase drowsiness
(Nighttime only)
■ avoid alcoholic beverages (Nighttime only)

Drug Facts (continued)

■ be careful when driving a motor vehicle or operating
machinery (Nighttime only)

Stop use and ask a doctor if

■ nervousness, dizziness, or sleeplessness occur
■ pain, nasal congestion, or cough gets worse or lasts more than
7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present ■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts.
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before
use.

Keep out of reach of children. In case of accidental overdose,
get medical help or contact a Poison Control Center
right away. Prompt medical attention is critical for adults
as well as for children even if you do not notice any signs
or symptoms.

If taking NIGHTTIME and DAYTIME products, carefully read
each section to ensure correct dosing. Do not take DAY &
NIGHT at the same time.

ADHESIVE AREA

ADHESIVE AREA

Drug Facts (continued)

Directions

■ do not use more than directed
■ do not take more than 12 caplets in any 24-hour period
■ adults and children 12 years and over: take 2 caplets every 4
hours
■ children under 12 years: do not use

Other information

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS
OPENED OR BLISTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C
(59°-86°F)
■ see end flap for expiration date and lot number

Inactive ingredients (Daytime only)

corn starch, croscarmellose sodium, croscopolone, FD&C red #40 aluminum lake, FD&C
yellow #6 aluminum lake, magnesium stearate, maltodextrin,
microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol,
povidone, silicon dioxide, sodium starch glycolate, stearic acid,
talc, titanium dioxide

Drug Facts (continued)

Inactive ingredients (Nighttime only)

corn starch, croscarmellose sodium, croscopolone, FD&C blue #1
aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow,
magnesium stearate, methacrylic acid and ethyl acrylate
copolymer, microcrystalline cellulose, polyethylene glycol,
polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate,
stearic acid, talc, titanium dioxide

Questions? Call 1-800-910-5874

094 03 0452 R00 C-001472-01-109
Distributed by Target Corporation
Minneapolis, MN 55403
©2021 Target Brands, Inc.

50844
REV0517B61569409
R-1920-615694-09SM-DF

ADHESIVE AREA

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

[illegible]

NDC 11673-336-06

Compare to the active ingredients in
Maximum Strength Mucinex® Fast-Max®
Cold, Flu & Sore Throat*

maximum strength
multi-symptom

fast mucus relief
cold, flu
and sore throat

acetaminophen

(pain reliever/fever reducer)

dextromethorphan HBr (cough suppressant)

guaifenesin (expectorant)

phenylephrine HCl (nasal decongestant)

relieves headache and fever

controls cough

relieves nasal and chest congestion

thins and loosens mucus



AGES
12+
YEARS

PLD-B282C
LB002840

094 14 0042 R01 ID305418
Dist. by Target Corp., Mpls., MN 55403

Product of U.S.A.

©2017 Target Brands, Inc.

Questions? Call 1-800-910-6874



TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY
SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt
Benckiser, distributor of Maximum Strength Mucinex®
Fast-Max® Cold, Flu & Sore Throat.

PLD-C282C LB003439

PARENTS:
Learn about how medicine abuse
www.StopMedicineAbuse.org

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses

- temporarily relieves these common cold and flu symptoms ■ sinus congestion and pressure
- minor aches and pains ■ nasal congestion
- cough due to minor throat and bronchial irritation ■ sore throat ■ headache
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash
- If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease ■ heart disease ■ diabetes
- thyroid disease ■ high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-800-910-6874

NDC 11673-337-06

Compare to active ingredients in **Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough***

maximum strength
fast mucus relief
severe congestion
and cough

dextromethorphan HBr 20 mg
(cough suppressant)
guaifenesin 400 mg (expectorant)
phenylephrine HCl 10 mg
(nasal decongestant)

controls cough
relieves nasal and chest congestion
thins and loosens mucus



AGES
12+
YEARS

6 FL OZ (177 mL)

PLD-B283D LB004222

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY
SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough.



094 14 0290 R01
C-000274-01-049
Dist. by Target Corp.
Minneapolis, MN 55403

TM & ©2021 Target Brands, Inc. PLD-B283D LB004223

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts

Active ingredients (in each 20 mL) Purposes

Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep
- nasal congestion due to a cold

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Drug Facts (continued)

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 17 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Drug Facts (continued)

Questions or comments?

Call 1-800-910-6874

PEEL CORNER FOR MORE DRUG FACTS

